

Content

In Brief

History and Current Situation

ENSO Outlook

Status of ENSO Indicators

Explanatory Note: El Niño and La Niña

In Brief

- *El-Niño continues to persist in the tropical Pacific Ocean and is nearing its end.*
- *Sea surface temperatures in the central tropical Pacific will continue to cool in the coming months.*
- *El Niño is likely to persist until the end of April or May 2024, with a transition to ENSO-neutral state likely during April to June 2024.*
- *Fiji usually experiences below normal rainfall during an El Niño event, however, impacts are likely to be witnessed during the onset of dry season.*
- *Fiji Met Service will continue to monitor the ENSO conditions closely and provide updates accordingly.*

History and Current Situation

History

The sea surface temperatures in the central and eastern equatorial Pacific Ocean warmed during July 2023, with most oceanic and atmospheric indicators implying an establishment of a weak El-Niño event. Since then the Pacific Ocean has been consistent with a weak El-Niño event. From October onwards, the event intensified into a moderate El Niño, peaked in December, and is now nearing its end.

Current Situation

El-Niño event continued to persist within the tropical Pacific Ocean. SSTs were above average across most the Pacific Ocean, with the largest anomalies in the central and east-central Pacific Ocean. Positive subsurface temperature anomalies have weakened across the equatorial Pacific, remaining close to the surface in the central Pacific. Negative subsurface temperature anomalies have expanded across the equatorial Pacific. Recently, below-average temperatures have reached the surface in the eastern Pacific Ocean.

The SOI for February 2024 was -12.6 , with the 5-month running mean of -5.3 . The latest 30-days average SOI until 22nd February 2024 was $+3.0$. Trade winds have been generally close to average over most of the equatorial Pacific. Cloudiness near the equatorial Date Line has been fluctuating around the average, but in the past two weeks, it has decreased. This is an indication that the presence of atmospheric El Niño conditions are weakening in the tropical Pacific. However, overall, the atmospheric and oceanic indicators are still consistent with the presence of an El Niño event.


ENSO Outlook

Sea surface temperatures in the central Pacific are likely to cool in the coming months, with a return to neutral ENSO levels expected around the April to June 2024 period.

The current El Niño event continues, nearing its end with a transition to an ENSO-neutral state likely during April to June 2024. FMS will continue to monitor the ENSO conditions closely and provide updates accordingly.

Fiji usually experiences *below normal* rainfall during an El Niño event, with the impacts more likely to be witnessed during the onset of the dry season.

Figure 1: Sea Surface Temperatures (SSTs) in the Pacific Ocean




SSTs were above average across most of the Pacific Ocean, with the largest anomalies in the central and east-central Pacific Ocean.

[Sustained warm SSTs in the equatorial Pacific Ocean are associated with El Niño events and cool anomalies with La Niña events].

Image source: USA's National Oceanic and Atmospheric Administration (NOAA).

Figure 2: Sub-surface Waters in the Equatorial Pacific Ocean




Positive subsurface temperature anomalies have weakened across the equatorial Pacific, remaining close to the surface in the central Pacific. Negative subsurface temperature anomalies have expanded across the equatorial Pacific. Recently, below-average temperatures have reached the surface in the eastern Pacific Ocean (near 120° - 90°W).

[Waters below the surface of the Ocean are good indicator of what may eventually happen at the surface in the coming months].

Image source: NOAA.

Most recent pentad analysis


Figure 3: Southern Oscillation Index (SOI)



The SOI for February 2024 was -12.6, with the 5-month running mean of -5.3. The latest 30-days average SOI to 22nd March 2024 was +3.0.

[Sustained values of SOI above +7 indicate presence of La Niña event and sustained values below -7 signify El Niño event].

Figure 4 : Near surface winds in the Pacific Ocean




Trade winds have been generally close to average over most of the equatorial Pacific.

[During El Niño there is a sustained weakening, or reversal, of the trade winds across much of the tropical Pacific. Conversely, during La Niña, there is a sustained strengthening of the trade winds].

Image source: NOAA.

Figure 5 : Cloudiness near the Dateline




Cloudiness near the equatorial Date Line has been fluctuating around average, but in the past two weeks, it has decreased. This suggests that the atmospheric effects of El Niño are weakening in the Pacific.

[Equatorial cloudiness near the Date Line typically increases during El Niño (negative OLR anomalies) and decreases during La Niña (positive OLR anomalies)].

Image source: Australian Bureau of Meteorology.

Figure 6: Climate Model Predictions of ENSO



Climate models on average show that the current El Niño will persist through March to May 2024 and then transition to ENSO-neutral state during April to June 2024.

Image source: International Research Institute for Climate and Society.

Explanatory Note - El Niño and La Niña

ENSO is an irregular cycle of persistent warming and cooling of SSTs in the tropical Pacific Ocean. The warm extreme is known as El Niño and cold extreme, La Niña.

The term El Niño was given to a warming of the ocean near the Peruvian coast in South America that appears around Christmas. Scientists now refer to an El Niño event as sustained warming over a large part of central and eastern equatorial Pacific Ocean. This warming is usually accompanied by persistent negative values of Southern Oscillation Index (SOI), a decrease in the strength or reversal of the Trade winds, increase in cloudiness near Dateline in the equatorial Pacific and a reduction in rainfall over most of Fiji (not immediate effect as there is a lag period) which can, especially during moderate to strong events, lead to drought.

La Niña is a sustained cooling of the central and eastern equatorial Pacific Ocean. The cooling is usually accompanied by persistent positive values of SOI, an increase in strength of the equatorial Trade winds, decrease in cloudiness near the Dateline in the equatorial Pacific and higher than average rainfall for most of Fiji (not immediate effects as there is a lag period), with frequent and sometimes severe flooding, especially during the wet season (November to April).