



Fiji Meteorological Service

ISO 9001:2015
certified Climate Services

Volume: 21 Issue: 4
Issued: March 31, 2025

Climate Outlook for Hydro-electricity Generation from April to June 2025

Current Conditions

Fiji's Climate

During 1st to 30th March, mainly fine weather prevailed, with some showers and thunderstorms, heavy at times, experienced over some parts of the country towards the end of the month.

There were 21 rainfall stations that reported in, in time for the compilation of this bulletin, with 2 stations reporting *well below average*, 13 *below average*, 5 *average* and 1 stations reporting *above average* rainfall.

The total monthly rainfall at Monasavu, until 30th March was 276mm, which is in *below average* category (55% of *normal*), when compared against the WMO standard 30-year average.

During January until 30th March 2025, Monasavu recorded 1244mm of rainfall, which was 74% of the *normal*, while in the past 6 months (October to 30th March, 3061mm of rainfall was recorded (97% of the *normal*) at the station (Figure 1).

El Niño Southern Oscillation (ENSO) Status

ENSO status is currently in neutral, with likely chances of the event to persist until August according to global climate models. Sea surface temperatures (SSTs) are currently *near to above average* in the eastern and far western Pacific and *below average* in the central Pacific ocean.

The Southern Oscillation Index (SOI) for February 2025 was 7.7, with the 5-month running mean of 6.6. The latest 30-day value to 23rd March 2025 was 10.0.

Trade winds have been closer to average in the western and the central Pacific. Cloudiness has been above average. Overall, ENSO indicators currently reflect neutral conditions, with the likelihood of persisting in the coming months.

El Niño-Southern Oscillation and Monasavu Climate Predictions

El-Niño Southern Oscillation Prediction

Recently surveyed global climate models, on average favor neutral conditions during the April to June period, which is likely to continue until August.

Minimum & Maximum Air Temperature Predictions - April & April to June 2025:

Day and night time temperatures are both likely to be *above normal* across Viti Levu and Vanua Levu during April as well as the April to June 2025 period (Figure 3).

Rainfall Predictions:

Fortnightly: 30th March– 12th April & 6th – 19th April

Rainfall across Viti Levu is likely to be above median from 30th March– 12th April, as well as from 6th to 19th April.

April 2025

There is 75% chance of receiving at least 148mm of rainfall at Nadarivatu station, 75% chance of at least 143mm of rainfall at the Nadarivatu and Monasavu

Dams and 75% chance of receiving at least 161mm of rainfall at Wailoa. There confidence in this forecast is Good (Table 1).

April to June 2025

For the April to June 2025 period, there is 75% chance of receiving at least 439mm of rainfall at Nadarivatu station, 75% chance of at least 483mm of rainfall at Nadarivatu Dam and Monasavu, and 75% chance of receiving at least 516mm of rainfall at Wailoa. There is high confidence on the generated outlook (Table 1).

Summary

There is no strong biasness for drier or wetter than usual condition across Viti Levu during April and April to June 2025 period.

Skill confidence is good to high for both the above forecasted periods.




Table 1: Rainfall Outlook: April & April to June 2025

April Outlook				
	25% chance of at least (mm)	50% chance of at least (mm)	75% chance of at least (mm)	Forecast Confidence
Nadarivatu station	388	310	148	Good
Nadarivatu Dam	397	300	143	Good
Monasavu Dam	397	300	143	Good
Wailoa	420	337	161	Good
April to June Outlook				
	25% chance of at least (mm)	50% chance of at least (mm)	75% chance of at least (mm)	Forecast Confidence
Nadarivatu station	805	643	439	High
Nadarivatu Dam	827	678	483	High
Monasavu Dam	827	678	483	High
Wailoa	853	696	516	High

The table above provides 25%, 50% and 75% chances of each station receiving the amount of rainfall mentioned above.

Figure 1: Rainfall Outlook: Fortnightly: 30th March – 12th April & 6th – 19th April




Figure 2: Rainfall Outlook: April & April to June 2025






Figure 3: Minimum Air Temperature Predictions: April & April to June 2025



Minimum air temperatures are expected to be *above normal* across Viti Levu and Vanua Levu, during April and April to June 2025 period. *Source: ACCESS-S2 Model.*

Figure 3: Maximum Air Temperature Predictions: April & April to June 2025



Maximum air temperatures are likely to be *above normal* across Viti Levu and Vanua Levu, during April and April to June 2025 period. *Source: ACCESS-S2 Model.*

Explanatory Notes

Climate Outlook for Hydro-electricity Generation is produced to provide advisories to Energy Fiji Limited (EFL). It aims to provide advanced warning on climate abnormalities for planning on economic generation mix and hydro-storage optimization.

Climate (Rainfall/Air Temperature) Outlook

Above normal – indicates that the rainfall/temperature value lies in the highest third of observation recorded in the standard 30 year normal period.

Near normal – indicates that the rainfall/temperature value lies in the middle third of observation recorded in the standard 30 year normal period.

Below normal – indicates that the rainfall/temperature value lies in the lowest third of observation recorded in the standard 30 year normal period.

Climatology – means that there are equal chances of receiving below normal, normal and above normal rainfall.

Median – rainfall value which marks the level dividing the ranked data set in half, that is, the midpoint of the ordered (lowest to highest) monthly or yearly rainfall totals.

Above Median – rainfall value that lies above the median value.

Below Median – rainfall value that lies below the median value.

El Niño Southern Oscillation (ENSO)

ENSO is the principal driver of the year-to-year variability of Fiji's climate. There are three phases of this phenomenon, *El Niño*, *La Niña* and *Neutral* conditions. El Niño or La Niña events are a natural part of the global climate system and usually recur after every 2 to 7 years. It normally develops around April to June, attains peak intensity between December to February and usually starts to decay around April to June period the following year. While most events last for a year, some have persisted for up to 2 years. It should be also noted that no two El Niño or La Niña events are the same. Different events have different impacts, but most exhibit some common climate characteristics.

Usually there is a lag effect on Fiji's climate with ENSO events, that is, once an El Niño or La Niña event is established in the tropical Pacific, it may take 2-6 months before its impact is seen on Fiji. Similarly, once an event finishes, it can take 2-6 months for climate to normalise.

El Niño events are associated with warming of the central and eastern tropical Pacific. El Niño events usually result in reduction of Fiji's rainfall. Often the whole of Fiji is affected in varying degrees and it is quite unusual for one part of the country to experience a prolonged dry spell, while the other is in a wet spell. The relationship and level of rainfall suppression is greater in the Dry Zone than in the Wet Zone. It is the suppression of rainfall during the Cool/Dry Season (May to October) that is normally of most concern. A reduction in Cool/Dry Season rainfall in the Dry Zone results in little or no rainfall until the next Wet Season. While usually the strength of an ENSO event is proportional to its impact on Fiji, at times weak event can also have a significant impact.

La Niña events are associated with cooling of the central and eastern tropical Pacific. Usually La Niña results in wetter than normal conditions for Fiji, occasionally leading to flooding during the Warm/Wet Season (November to April).

During **Neutral** condition, neither El Niño nor La Niña is present, it has little effect on global climate, meaning other climate influences are more likely to dominate.

Lag effects – means that there is a delay in a change of some aspect of climate due to influence of other factors that is acting slowly.

Climate bulletins that can be viewed together with this bulletin include:

- 1) *Fiji Climate Summary* at <https://www.met.gov.fj/index.php?page=FijiClimateSummary> (issued monthly)
- 2) *Fiji Climate Outlook* at <https://www.met.gov.fj/index.php?page=ClimateOutlook> (issued monthly)

This information is prepared as soon as ENSO, climate and oceanographic data is received from recording stations around Fiji and Meteorological Agencies around the world. While every effort is made to verify observational data, Fiji Meteorological Service does not guarantee the accuracy and reliability of the analyses presented, and accepts no liability for any losses incurred through the use of this information and its contents. The information may be freely disseminated provided the source is acknowledged. For further clarification and expert advice, please contact the Fiji Meteorological Service HQ, Namaka, Nadi.

For further information, contact: The Director of Meteorology, Fiji Meteorological Service, Private Mail Bag NAP0351, Nadi Airport, Fiji. Phone: (679) 6724888, Fax: (679) 6720430, E-mail: fms@met.gov.fj or climate@met.gov.fj. URL: <http://www.met.gov.fj>